

ON THE BETTI NUMBERS OF SHIFTED COMPLEXES OF STABLE SIMPLICIAL COMPLEXES

ZHONGMING TANG AND GUIFEN ZHUANG

ABSTRACT. Let Δ be a stable simplicial complex on n vertexes. Over an arbitrary base field K , the symmetric algebraic shifted complex Δ^s of Δ is defined. It is proved that the Betti numbers of the Stanley-Reisner ideals in the polynomial ring $K[x_1, x_2, \dots, x_n]$ of the symmetric algebraic shifted, exterior algebraic shifted and combinatorial shifted complexes of Δ are equal.

INTRODUCTION

Throughout the paper, let Δ be a simplicial complex on the vertex set $[n] = \{1, 2, \dots, n\}$. Then, by definition, Δ is a set of some subsets of $[n]$ such that $\{i\} \in \Delta$, $i = 1, 2, \dots, n$, and, for any $\sigma \in \Delta$, if $\tau \subseteq \sigma$ then $\tau \in \Delta$. Set $\nabla = \{\sigma \subseteq [n] : \sigma \notin \Delta\}$. Let $S = K[x_1, x_2, \dots, x_n]$ be the polynomial ring over an infinite field K . For any $\sigma = \{i_1, i_2, \dots, i_r\} \subseteq [n]$ with $i_1 < i_2 < \dots < i_r$, we set $x_\sigma = x_{i_1}x_{i_2} \cdots x_{i_r}$. Then Δ determines an ideal I_Δ , the Stanley-Reisner ideal of Δ , of S , which is the monomial ideal generated by all monomials x_σ such that $\sigma \in \nabla$. Conversely, for any monomial ideal L of S which is generated by some squarefree monomials, there is a unique simplicial complex Δ_1 on the vertex set $[n]$ such that $L = I_{\Delta_1}$. We refer the reader to [5] and [10] for the detailed information about these notions.

For any graded ideal I of S , the graded Betti number $\beta_{i,j}(I)$ is defined by the minimal graded free S -resolution of I :

$$\cdots \rightarrow \bigoplus_j S(-j)^{\beta_{i,j}(I)} \rightarrow \cdots \rightarrow \bigoplus_j S(-j)^{\beta_{1,j}(I)} \rightarrow \bigoplus_j S(-j)^{\beta_{0,j}(I)} \rightarrow I \rightarrow 0.$$

Set $\beta_i(I) = \sum_j \beta_{i,j}(I)$. The regularity, $\text{reg}(I)$, of I is defined as $\text{reg}(I) = \max\{j - i : \beta_{i,j}(I) \neq 0\}$.

For the simplicial complex Δ , there are three shifted complexes Δ^s , Δ^e and Δ^c , which are called the symmetric algebraic shifted complex, exterior algebraic shifted complex and combinatorial shifted complex of Δ respectively, cf. [1]. Shifted complexes are determined by some monomial ideals of a polynomial ring or an exterior algebra over a field K . The combinatorial shifting and the exterior algebraic shifting are defined over any base field K , while the symmetric algebraic shifting is only defined when $\text{char}(K) = 0$, cf. [1]. In the present paper, we firstly show that if I is a (squarefree) stable ideal of S , then $\text{Gin}(I)$ is also stable. Then it makes sense to define the symmetric algebraic shifting of stable complexes over an arbitrary base field K . One of the important problems in the study of simplicial complexes is the

Key words and phrases. shifted complexes, Stanley-Reisner ideals, Betti numbers.

2000 AMS Subject Classifications: 13F55, 13D02, 55U10

Supported by the National Natural Science Foundation of China.

behaviour of the graded Betti numbers under shifting. In [1], Aramova, Herzog and Hibi conjectured that $\beta_{i,j}(I_{\Delta^s}) \leq \beta_{i,j}(I_{\Delta^e}) \leq \beta_{i,j}(I_{\Delta^c})$, although their symmetric algebraic shifting is only defined in characteristic 0. In this paper, we will show that, over any base field, when Δ is stable, $\beta_{i,j}(I_{\Delta^s}) = \beta_{i,j}(I_{\Delta^e}) = \beta_{i,j}(I_{\Delta^c}) = \beta_{i,j}(I_{\Delta})$, for all i and j .

1. BETTI NUMBERS UNDER SYMMETRIC ALGEBRAIC SHIFTING

A monomial $x_1^{a_1} x_2^{a_2} \cdots x_n^{a_n} \in S$ is said to be squarefree if $a_i \leq 1$, $i = 1, 2, \dots, n$. For a monomial $u \in S$, we denote $\max\{i : x_i \mid u\}$ by $m(u)$, and, for any $\sigma = \{i_1, i_2, \dots, i_r\} \subseteq [n]$, we also denote $\max\{i_s : s = 1, 2, \dots, r\}$ by $m(\sigma)$. Let I be a monomial ideal of S , the minimal system of monomial generators of I will be denoted by $G(I)$. For any j , we write I_j ($G(I)_j$) for the set of monomials of degree j belonging to I ($G(I)$) and $I_{\langle j \rangle}$ for the ideal generated by I_j . A monomial ideal I of S is said to be squarefree if it is generated by squarefree monomials. Then, it is clear that I_{Δ} is a squarefree monomial ideal.

A (squarefree) monomial ideal is said to be (squarefree) stable if its set B of (squarefree) monomials is (squarefree) stable, i.e., $x_i(u/x_{m(u)}) \in B$ for all $u \in B$ and all $i < m(u)$ (such that $x_i \nmid u$), further, if $x_i(u/x_j) \in B$ for all $u \in B$ and all $i < j$ such that $x_j \mid u$ (and $x_i \nmid u$) then we say that the ideal is strongly (squarefree) stable. Let I be a monomial ideal of S . If I is stable, it is shown in [8] that

$$\beta_{i,i+j}(I) = \sum_{u \in G(I)_j} \binom{m(u) - 1}{i}$$

and

$$\text{reg}(I) = \max\{\deg(u) : u \in G(I)\}.$$

If I is squarefree stable, then it is given in [3] that

$$\beta_{i,i+j}(I) = \sum_{u \in G(I)_j} \binom{m(u) - j}{i}$$

and

$$\text{reg}(I) = \max\{\deg(u) : u \in G(I)\}.$$

We say that the simplicial complex Δ is stable if I_{Δ} is squarefree stable. Then, it is easy to show the following

Lemma 1.1. *The following are equivalent*

- (1) Δ is stable.
- (2) For any $\sigma \in \nabla$, if $1 \leq i < m(\sigma)$ and $i \notin \sigma$, then $(\sigma \setminus \{m(\sigma)\}) \cup \{i\} \in \nabla$.
- (3) For any $\sigma \in \Delta$, if $i \in \sigma$ and $m(\sigma) < j \leq n$, then $(\sigma \setminus \{i\}) \cup \{j\} \in \Delta$.

For a graded ideal I of S , let $\text{Gin}(I)$ be the generic initial ideal of I with respect to the reverse lexicographic term order on S induced by $x_1 > x_2 > \dots > x_n$. Then $\text{Gin}(I)$ is Borel-fixed and, when $\text{char}(K) = 0$, it is strongly stable, cf. [6, Chapter 15]. For a general field K , we have the following

Proposition 1.2. *If I is a (squarefree) stable ideal of S , then $\text{Gin}(I)$ is also stable.*

Proof. Let $j \geq 1$ be such that $I_j \neq 0$. As I is stable or squarefree stable, it follows that I is componentwise linear, hence $I_{\langle j \rangle}$ is j -linear. Note that $\text{reg}(I_{\langle j \rangle}) = j$. Then $\text{reg}(\text{Gin}(I_{\langle j \rangle})) = \text{reg}(I_{\langle j \rangle}) = j$, thus $\text{Gin}(I_{\langle j \rangle})$ is generated in degree j . By the construction of generic initial ideals, we see that $\text{Gin}(I)_j = \text{Gin}(I_{\langle j \rangle})_j$, then it turns out that $\text{Gin}(I)_{\langle j \rangle} = \text{Gin}(I_{\langle j \rangle})$. Since $\text{Gin}(I_{\langle j \rangle})$ is Borel-fixed and generated in degree j and $\text{reg}(\text{Gin}(I_{\langle j \rangle})) = j$, it follows from [7, Proposition 10] that $\text{Gin}(I_{\langle j \rangle})$ is stable, i.e., $\text{Gin}(I)_{\langle j \rangle}$ is stable. Hence $\text{Gin}(I)$ is stable. $\square$

There is an operator σ which transfers any monomial ideal into a squarefree monomial ideal. For any $u = x_{i_1}x_{i_2} \cdots x_{i_j} \cdots x_{i_d} \in S$ where $i_1 \leq i_2 \leq \cdots \leq i_j \leq \cdots \leq i_d$, we set $u^\sigma = x_{i_1}x_{i_2+1} \cdots x_{i_j+(j-1)} \cdots x_{i_d+(d-1)}$. Let I be a monomial ideal of S with $G(I) = \{u_1, u_2, \dots, u_r\}$. Then we set I^σ for the squarefree monomial ideal generated by $u_1^\sigma, u_2^\sigma, \dots, u_r^\sigma$. Thus I^σ is a squarefree ideal of $K[x_1, x_2, \dots, x_m]$, where $m = \max\{m(u) + \deg(u) - 1 : u \in G(I)\}$.

Suppose that Δ is stable. Then, by 1.2, $\text{Gin}(I_\Delta)$ is also stable. Using the same arguments as in the proof of [1, Lemma 1.1], we see that $m(u) + \deg(u) - 1 \leq n$ for all $u \in G(\text{Gin}(I_\Delta))$, hence, $(\text{Gin}(I_\Delta))^\sigma$ is also an ideal of S . Then we define the symmetric algebraic shifted complex Δ^s of Δ by

$$I_{\Delta^s} = (\text{Gin}(I_\Delta))^\sigma.$$

We are going to show that $\beta_{i,j}(I_\Delta) = \beta_{i,j}(I_{\Delta^s})$ if Δ is stable. When $\text{char}(K) = 0$, this result can be easily proved. But, for a base field K with arbitrary characteristic, we need some new results as we do not know whether $\text{Gin}(I_\Delta)$ is strongly stable.

Proposition 1.3. *If I is a (squarefree) stable ideal of S , then*

$$\beta_{i,j}(I) = \beta_{i,j}(\text{Gin}(I)) \text{ for all } i \text{ and } j,$$

Proof. By 1.2, $\text{Gin}(I)$ is also stable, thus I and $\text{Gin}(I)$ are componentwise linear. Then, by [9, Proposition 1.3], for any i and j ,

$$\begin{aligned} \beta_{i,i+j}(I) &= \beta_i(I_{\langle j \rangle}) - \beta_i(\mathbf{m}I_{\langle j-1 \rangle}), \\ \beta_{i,i+j}(\text{Gin}(I)) &= \beta_i(\text{Gin}(I)_{\langle j \rangle}) - \beta_i(\mathbf{m}\text{Gin}(I)_{\langle j-1 \rangle}) \\ &= \beta_i(\text{Gin}(I_{\langle j \rangle})) - \beta_i(\mathbf{m}\text{Gin}(I_{\langle j-1 \rangle})), \end{aligned}$$

where $\mathbf{m}$ is the ideal $(x_1, x_2, \dots, x_n)$ of S . Since $\mathbf{m}I_{\langle j-1 \rangle}$ is j -linear, it follows that $\text{reg}(\mathbf{m}I_{\langle j-1 \rangle}) = j$, hence $\text{reg}(\text{Gin}(\mathbf{m}I_{\langle j-1 \rangle})) = j$. Then $\text{Gin}(\mathbf{m}I_{\langle j-1 \rangle})$ is also generated in degree j . On the other hand, as $\text{Gin}(I_{\langle j-1 \rangle})$ is generated in degree $j-1$ (refer to the proof of 1.2), we see that $\mathbf{m}\text{Gin}(I_{\langle j-1 \rangle})$ is also generated in degree j . But

$$\text{Gin}(\mathbf{m}I_{\langle j-1 \rangle})_j = \text{Gin}(I_{\langle j-1 \rangle})_j = (\mathbf{m}\text{Gin}(I_{\langle j-1 \rangle}))_j,$$

hence $\mathbf{m}\text{Gin}(I_{\langle j-1 \rangle}) = \text{Gin}(\mathbf{m}I_{\langle j-1 \rangle})$. To end the proof, it is enough to show that, if I is (squarefree) stable and generated in degree d , then $\beta_i(I) = \beta_i(\text{Gin}(I))$. By the assumptions, I and $\text{Gin}(I)$ are d -linear and their Hilbert functions have the following

forms:

$$H_I(t) = \left(\sum_i (-1)^i \beta_{i,i+d}(I) t^i \right) \frac{t^d}{(1-t)^n},$$

$$H_{\text{Gin}(I)}(t) = \left(\sum_i (-1)^i \beta_{i,i+d}(\text{Gin}(I)) t^i \right) \frac{t^d}{(1-t)^n}.$$

But I and $\text{Gin}(I)$ have the same Hilbert function, hence $\beta_i(I) = \beta_i(\text{Gin}(I))$, as required. $\square$

The following result is proved in [1] when I is strongly stable.

Lemma 1.4. *If I is stable and Borel-fixed, then $\beta_{i,j}(I) = \beta_{i,j}(I^\sigma)$, for all i and j .*

Proof. Let $G(I) = \{u_1, u_2, \dots, u_r\}$. We will show that I^σ is squarefree stable with $G(I^\sigma) = \{u_1^\sigma, u_2^\sigma, \dots, u_r^\sigma\}$. Then the result follows from the formulas of Betti numbers of stable and squarefree stable ideals.

Let $u = x_{i_1}^{r_1} x_{i_2}^{r_2} \cdots x_{i_s}^{r_s} \in I$ with $i_1 < i_2 < \dots < i_s$ and $r_j \geq 1$, $j = 1, 2, \dots, s$. We firstly note that, since I is p -Borel with $p = \text{char}(K)$ and $k \leq_p k$ for any positive integer k , cf. [6, Chapter 15], it follows that $(x_k/x_{i_j})^{r_j} u \in I$ for all $1 \leq k < i_j$. Secondly, let $v = x_{j_1}^{t_1} x_{j_2}^{t_2} \cdots x_{j_l}^{t_l} \in I$ with $j_1 < j_2 < \dots < j_l$ and $t_k \geq 1$, $k = 1, 2, \dots, l$. Suppose that $v^\sigma \mid u^\sigma$ and to see the relationship between u and v . For any $r \in [n]$ and $J \subseteq [n] \cup \{0\}$, set $x_{r+J} = \prod_{k \in J} x_{r+k}$ and $[0, r-1] = \{0, 1, \dots, r-1\}$. Then

$$u^\sigma = x_{i_1+[0,r_1-1]} \cdot x_{i_2+r_1+[0,r_2-1]} \cdots x_{i_s+(r_1+\dots+r_{s-1})+[0,r_s-1]}.$$

We notice that the difference between indices of any two x_w in different groups $x_{i_k+(r_1+\dots+r_{k-1})+[0,r_k-1]}$ is at least 2. Thus, from

$$v^\sigma = x_{j_1+[0,t_1-1]} \cdot x_{j_2+t_1+[0,t_2-1]} \cdots x_{j_l+(t_1+\dots+t_{l-1})+[0,t_l-1]}$$

and $v^\sigma \mid u^\sigma$, we see that $l \leq s$ and there exist $w_k \in [s]$, $k = 1, 2, \dots, l$, such that

$$x_{j_k+(t_1+\dots+t_{k-1})+[0,t_k-1]} \mid x_{i_{w_k}+(r_1+\dots+r_{w_k-1})+[0,r_{w_k}-1]}, k = 1, 2, \dots, l.$$

Hence $j_k = i_{w_k} + m_k$ with $m_k \geq 0$ and $t_k \leq r_{w_k}$, $k = 1, 2, \dots, l$. It turns out that $x_{j_1-m_1}^{t_1} x_{j_2-m_2}^{t_2} \cdots x_{j_l-m_l}^{t_l} \mid u$ for some $m_k \geq 0$, $k = 1, 2, \dots, l$.

Now we show that $G(I^\sigma) = \{u_1^\sigma, u_2^\sigma, \dots, u_r^\sigma\}$. Suppose that $u_k^\sigma \notin G(I^\sigma)$. Then there exists some $l \neq k$ such that $u_l^\sigma \mid u_k^\sigma$. Let $u_l = x_{i_1}^{r_1} x_{i_2}^{r_2} \cdots x_{i_s}^{r_s}$ with $i_1 < i_2 < \dots < i_s$ and $r_j \geq 1$, $j = 1, 2, \dots, s$. Then $x_{i_1-m_1}^{r_1} x_{i_2-m_2}^{r_2} \cdots x_{i_s-m_s}^{r_s} \mid u_k$ for some $m_t \geq 0$, $t = 1, 2, \dots, s$. Note that $x_{i_1-m_1}^{r_1} x_{i_2-m_2}^{r_2} \cdots x_{i_s-m_s}^{r_s} \neq u_k$ by $u_l^\sigma \mid u_k^\sigma$. Since $x_{i_1}^{r_1} x_{i_2}^{r_2} \cdots x_{i_s}^{r_s} \in I$ and I is Borel-fixed, as noted above, we see that $x_{i_1-m_1}^{r_1} x_{i_2-m_2}^{r_2} \cdots x_{i_s-m_s}^{r_s} \in I$. This contradicts that $u_k \in G(I)$. Thus $G(I^\sigma) = \{u_1^\sigma, u_2^\sigma, \dots, u_r^\sigma\}$.

Finally, we show that I^σ is squarefree stable. Let $u = x_{i_1}^{r_1} x_{i_2}^{r_2} \cdots x_{i_s}^{r_s} \in G(I)$ with $i_1 < i_2 < \dots < i_s$ and $r_j \geq 1$, $j = 1, 2, \dots, s$. Suppose that $l < m(u^\sigma) = i_s + (r_1 + \dots + r_s) - 1$ and $x_l \nmid u^\sigma$. Let $i_{k-1} + (r_1 + \dots + r_{k-1}) - 1 < l < i_k + (r_1 + \dots + r_{k-1})$ and set $v = x_{i_1}^{r_1} \cdots x_{i_{k-1}}^{r_{k-1}} x_{l-(r_1+\dots+r_{k-1})} x_{i_k-1}^{r_k} \cdots x_{i_{s-1}-1}^{r_{s-1}} x_{i_s-1}^{r_s}$. Note that $v \in I$ as above and $v^\sigma = (x_l/x_{m(u^\sigma)}) u^\sigma$. Then $v = u_w v^*$ for some $u_w \in G(I)$ and a monomial $v^* \in S$, as I is stable, we may assume that $m(u_w) < j$ for all j such that $x_j \mid v^*$. Thus we have that $u_w^\sigma \mid v^\sigma$, hence $v^\sigma \in I^\sigma$, as required. $\square$

Now we prove the following

Theorem 1.5. *If Δ is stable, then, for all i and j ,*

$$\beta_{i,j}(I_\Delta) = \beta_{i,j}(I_{\Delta^s}).$$

Proof. Since I_Δ is squarefree stable, it follows from 1.3 that $\beta_{i,j}(I_\Delta) = \beta_{i,j}(\text{Gin}(I_\Delta))$. But $\text{Gin}(I_\Delta)$ is also stable by 1.2, and Borel-fixed. Then the theorem follows from 1.4 . $\square$

For exterior algebraic shifting, a similar result can be easily obtained. Let V be an n -dimensional K -vector space with basis $e_1, e_2, \dots, e_n$. Then the exterior algebra $E = \bigwedge(V)$ is a finite dimensional graded K -algebra and its r -th graded component $\bigwedge^r(V)$ has the K -basis $e_{i_1} \wedge e_{i_2} \wedge \dots \wedge e_{i_r}$ with $i_1 < i_2 < \dots < i_r$. For any $\sigma = \{i_1, i_2, \dots, i_r\} \subseteq [n]$ with $i_1 < i_2 < \dots < i_r$, we set $e_\sigma = e_{i_1} \wedge e_{i_2} \wedge \dots \wedge e_{i_r}$. The stable and strongly stable ideals of E (which are automatically squarefree) can be defined as in S . Let I be a graded ideal of E . The generic initial ideal, $\text{Gin}(I)$, of I in E is also defined similarly as in S . Then $\text{Gin}(I)$ is strongly stable for any infinite field K , cf. [4].

For the simplicial complex Δ , we set J_Δ , the Stanley-Reisner ideal of Δ in E , as the monomial ideal of E generated by all monomials e_σ such that $\sigma \in \nabla$. The exterior algebraic shifted complex Δ^e of Δ is defined by

$$J_{\Delta^e} = \text{Gin}(J_\Delta).$$

Suppose that Δ is stable. Then J_Δ is also componentwise linear. Thus, by [2, Corollary 2.2], we have the following

Proposition 1.6. *If Δ is stable, then, for all i and j ,*

$$\beta_{i,j}(I_\Delta) = \beta_{i,j}(I_{\Delta^e}).$$

2. BETTI NUMBERS UNDER COMBINATORIAL SHIFTING

Let $1 \leq k < l \leq n$, for any $\sigma \in \nabla$, we define

$$S_{kl}(\sigma) = \begin{cases} (\sigma \setminus \{l\}) \cup \{k\} & \text{if } l \in \sigma, k \notin \sigma \text{ and } (\sigma \setminus \{l\}) \cup \{k\} \notin \nabla, \\ \sigma & \text{otherwise.} \end{cases}$$

We summarize some properties of S_{kl} as follows:

- (1) For any $\sigma \in \nabla$, $S_{kl}(\sigma) \notin \nabla$ if $S_{kl}(\sigma) \neq \sigma$.
- (2) Let $\sigma, \tau \in \nabla$, if $\sigma \neq \tau$ then $S_{kl}(\sigma) \neq S_{kl}(\tau)$.
- (3) If Δ is stable, then $m(S_{kl}(\sigma)) = m(\sigma)$.
- (4) Let $G(I_\Delta) = \{x_{\sigma_1}, x_{\sigma_2}, \dots, x_{\sigma_r}\}$. Then $\sigma_i \neq S_{kl}(\sigma_j)$ for any $i \neq j$, and, if $S_{kl}(\sigma_i) \subseteq S_{kl}(\sigma_j)$ for $i \neq j$, then $S_{kl}(\sigma_i) \neq \sigma_i$ and $S_{kl}(\sigma_j) = \sigma_j$.

$\text{Shift}_{kl}(\Delta)$ is the simplicial complex defined by $I_{\text{Shift}_{kl}(\Delta)}$ which is the squarefree monomial ideal of S generated by all monomials $x_{S_{kl}(\sigma)}$ with $\sigma \in \nabla$. A combinatorial shifted complex Δ^c of Δ is a shifted complex and

$$\Delta^c = \text{Shift}_{k_r l_r}(\text{Shift}_{k_{r-1} l_{r-1}}(\dots (\text{Shift}_{k_1 l_1}(\Delta)) \dots)),$$

for some $r \geq 1$ and $k_i < l_i$, $i = 1, 2, \dots, r$.

We firstly discuss the properties of $\text{Shift}_{kl}(\Delta)$ and, then, get the properties of Δ^c by induction.

Lemma 2.1. *If Δ is stable, then $\text{Shift}_{kl}(\Delta)$ is also stable.*

Proof. It is enough to show that, for any $\sigma \in \nabla$, if $i < m(S_{kl}(\sigma))$ and $i \notin S_{kl}(\sigma)$, then $(S_{kl}(\sigma) \setminus \{m(S_{kl}(\sigma))\}) \cup \{i\} = S_{kl}(\sigma')$ for some $\sigma' \in \nabla$. Set $\sigma_i = (S_{kl}(\sigma) \setminus \{m(S_{kl}(\sigma))\}) \cup \{i\}$. Consider two cases.

Case I) $S_{kl}(\sigma) = \sigma$. Then $\sigma_i = (\sigma \setminus \{m(\sigma)\}) \cup \{i\} \in \nabla$ by the stability of Δ . We will show that $\sigma_i = S_{kl}(\sigma_i)$. On the contrary, suppose that $S_{kl}(\sigma_i) \neq \sigma_i$. Then $S_{kl}(\sigma_i) = (\sigma_i \setminus \{l\}) \cup \{k\}$, $l \in \sigma_i$, $k \notin \sigma_i$ and $(\sigma_i \setminus \{l\}) \cup \{k\} \notin \nabla$. Note that $i \neq k$ and $(\sigma_i \setminus \{l\}) \cup \{k\} = (((\sigma \setminus \{m(\sigma)\}) \cup \{i\}) \setminus \{l\}) \cup \{k\}$. Consider two possibilities: $i = l$ and $i \neq l$. When $i = l$, as Δ is stable, $k < m(\sigma)$ and $k \notin \sigma$, we have that

$$(\sigma_i \setminus \{l\}) \cup \{k\} = (\sigma \setminus \{m(\sigma)\}) \cup \{k\} \in \nabla,$$

a contradiction. When $i \neq l$, as $l \in \sigma_i$, $k \notin \sigma_i$ and $k < l$, we see that $l \in \sigma$ and $k \notin \sigma$. But $S_{kl}(\sigma) = \sigma$, this implies that $(\sigma \setminus \{l\}) \cup \{k\} \in \nabla$. By $l \in \sigma_i$ again, we have that $l \neq m(\sigma)$, hence $m((\sigma \setminus \{l\}) \cup \{k\}) = m(\sigma)$ and

$$(\sigma_i \setminus \{l\}) \cup \{k\} = (((\sigma \setminus \{l\}) \cup \{k\}) \setminus \{m((\sigma \setminus \{l\}) \cup \{k\})\}) \cup \{i\}.$$

Since $i < m(\sigma)$ and $i \notin (\sigma \setminus \{l\}) \cup \{k\}$, it follows from the stability of Δ that the right-hand side of the above equality is in ∇ , i.e., $(\sigma_i \setminus \{l\}) \cup \{k\} \in \nabla$, a contradiction again.

Case II) $S_{kl}(\sigma) \neq \sigma$. Then $S_{kl}(\sigma) = (\sigma \setminus \{l\}) \cup \{k\}$, $l \in \sigma$, $k \notin \sigma$ and $(\sigma \setminus \{l\}) \cup \{k\} \notin \nabla$. Thus, by the stability of Δ , we see immediately that $l \neq m(\sigma)$, hence $m(S_{kl}(\sigma)) = m(\sigma)$ and $\sigma_i = (((\sigma \setminus \{l\}) \cup \{k\}) \setminus \{m(\sigma)\}) \cup \{i\}$. Note that $i \neq k$ by $i \notin S_{kl}(\sigma)$. Consider two possibilities: $i = l$ and $i \neq l$ again. When $i = l$, as $k < m(\sigma)$ and $k \notin \sigma$, we have that $\sigma_i = (\sigma \setminus \{m(\sigma)\}) \cup \{k\} \in \nabla$ by the stability of Δ , and $S_{kl}(\sigma_i) = \sigma_i$, as required. When $i \neq l$, we have that $\sigma_i = (((\sigma \setminus \{m(\sigma)\}) \cup \{i\}) \setminus \{l\}) \cup \{k\}$. As $i < m(\sigma)$ and $i \notin \sigma$, we see that $(\sigma \setminus \{m(\sigma)\}) \cup \{i\} \in \nabla$ by the stability of Δ . Since $l \in (\sigma \setminus \{m(\sigma)\}) \cup \{i\}$ and $k \notin (\sigma \setminus \{m(\sigma)\}) \cup \{i\}$ by $k \notin \sigma$ and $i \neq k$, it follows that $\sigma_i = S_{kl}(\sigma_i)$ if $\sigma_i \in \nabla$ and $\sigma_i = S_{kl}((\sigma \setminus \{m(\sigma)\}) \cup \{i\})$ if $\sigma_i \notin \nabla$, as required. $\square$

Suppose that Δ is stable and $G(I_\Delta) = \{x_{\sigma_1}, x_{\sigma_2}, \dots, x_{\sigma_r}\}$. For any $\sigma \in \nabla$, there is an $i \in [r]$ such that $\sigma = \sigma_i \cup \tau$ with $\tau \subseteq [n]$, where we understand always that $\sigma \cap \tau = \emptyset$. Let $\tau_1, \tau_2 \subseteq [n]$, we write $\tau_1 < \tau_2$ if $s < t$ for all $s \in \tau_1$ and $t \in \tau_2$. Note that, as Δ is stable, we can find some $j \in [r]$ such that $\sigma = \sigma_j \cup \tau'$ where $\tau' \subseteq [n]$ and $\sigma_j < \tau'$, and such a representation is unique.

Let $i \in [n]$. If $S_{kl}(\sigma_i) \supseteq S_{kl}(\sigma_j)$ for some $j \neq i$, then $S_{kl}(\sigma_i) = \sigma_i$ and $S_{kl}(\sigma_j) \neq \sigma_j$, hence $\sigma_i = S_{kl}(\sigma_j) \cup \tau$ with $\tau \subseteq [n]$ and $\tau \neq \emptyset$, and conversely, this equality implies that $S_{kl}(\sigma_j) \neq \sigma_j$, $k \in \sigma_i$, hence $S_{kl}(\sigma_i) = \sigma_i$ and $S_{kl}(\sigma_i) \supseteq S_{kl}(\sigma_j)$. Furthermore, $\sigma_j \cup \tau$ is uniquely defined by σ_i since, if $\sigma_i = S_{kl}(\sigma_{j'}) \cup \tau'$ with $j' \in [r]$ and $\tau' \subseteq [n]$ then $l \in \sigma_j, \sigma_{j'}$, $k \notin \sigma_j, \sigma_{j'}$ and $((\sigma_j \setminus \{l\}) \cup \{k\}) \cup \tau = ((\sigma_{j'} \setminus \{l\}) \cup \{k\}) \cup \tau'$, hence $\sigma_j \cup \tau = \sigma_{j'} \cup \tau'$ recalling that, by our convention, $S_{kl}(\sigma_j) \cap \tau = S_{kl}(\sigma_{j'}) \cap \tau' = \emptyset$. As noted above, we may assume that $\sigma_j < \tau$. Then it makes sense to give the following definition.

Let $i \in [r]$, we define σ_i^* as follows

$$\sigma_i^* = \begin{cases} \sigma_s \cup \tau & \text{if } \sigma_i = S_{kl}(\sigma_s) \cup \tau \text{ for some } s \neq i \\ S_{kl}(\sigma_i) & \text{otherwise} \end{cases}$$

Thus there are three possible values for σ_i^* . If $S_{kl}(\sigma_i) \neq \sigma_i$ then $\sigma_i^* = S_{kl}(\sigma_i) \notin \nabla$; if $S_{kl}(\sigma_i) = \sigma_i$ then $\sigma_i^* = \sigma_s \cup \tau$ or σ_i depending on whether $\sigma_i = S_{kl}(\sigma_s) \cup \tau$ for some $s \neq i$ or not. Note that $m(\sigma_i^*) = m(\sigma_i)$, $i = 1, 2, \dots, r$.

Lemma 2.2. *Suppose that Δ is stable. If $G(I_\Delta) = \{x_{\sigma_1}, x_{\sigma_2}, \dots, x_{\sigma_r}\}$, then*

$$G(I_{\text{Shift}_{kl}(\Delta)}) = \{x_{\sigma_1^*}, x_{\sigma_2^*}, \dots, x_{\sigma_r^*}\}.$$

Proof. Firstly, we show that $I_{\text{Shift}_{kl}(\Delta)}$ is generated by $x_{\sigma_1^*}, x_{\sigma_2^*}, \dots, x_{\sigma_r^*}$. It is enough to show that, for any $\sigma \in \nabla$, there is an i such that $S_{kl}(\sigma) = \sigma_i^* \cup \tau$. As $G(I_\Delta) = \{x_{\sigma_1}, x_{\sigma_2}, \dots, x_{\sigma_r}\}$, we have that $\sigma = \sigma_j \cup \tau_1$, for some j , where $\sigma_j < \tau_1$. Divide into three cases.

Case I) $S_{kl}(\sigma_j) \neq \sigma_j$. Then $\sigma_j^* = S_{kl}(\sigma_j)$, $l \in \sigma_j$, $k \notin \sigma_j$ and $S_{kl}(\sigma_j) = (\sigma_j \setminus \{l\}) \cup \{k\} \notin \nabla$, hence $l \in \sigma$. If $k \in \sigma$, then $k \in \tau_1$ and

$$\begin{aligned} S_{kl}(\sigma) &= \sigma \\ &= \sigma_j \cup \tau_1 \\ &= ((\sigma_j \setminus \{l\}) \cup \{k\}) \cup ((\tau_1 \setminus \{k\}) \cup \{l\}) \\ &= \sigma_j^* \cup ((\tau_1 \setminus \{k\}) \cup \{l\}), \end{aligned}$$

as required. If $k \notin \sigma$ and $(\sigma \setminus \{l\}) \cup \{k\} \in \nabla$, then $(\sigma \setminus \{l\}) \cup \{k\} = \sigma_t \cup \tau'$ for some t , where $\sigma_t < \tau'$, and $S_{kl}(\sigma) = \sigma$. Note that $(\sigma \setminus \{l\}) \cup \{k\} = ((\sigma_j \setminus \{l\}) \cup \{k\}) \cup \tau_1$. Then

$$((\sigma_j \setminus \{l\}) \cup \{k\}) \cup \tau_1 = \sigma_t \cup \tau',$$

where $(\sigma_j \setminus \{l\}) \cup \{k\} < \tau_1$ and $\sigma_t < \tau'$. As $(\sigma_j \setminus \{l\}) \cup \{k\} \notin \nabla$, we see that $(\sigma_j \setminus \{l\}) \cup \{k\} \subseteq \sigma_t$. Thus $\sigma_t = ((\sigma_j \setminus \{l\}) \cup \{k\}) \cup \tau'' = S_{kl}(\sigma_j) \cup \tau''$ with $\tau'' \subseteq \tau_1$, then $\sigma_t^* = \sigma_j \cup \tau''$. Hence

$$\begin{aligned} S_{kl}(\sigma) &= \sigma \\ &= \sigma_j \cup \tau_1 \\ &= (\sigma_j \cup \tau'') \cup (\tau_1 \setminus \tau'') \\ &= \sigma_t^* \cup (\tau_1 \setminus \tau''), \end{aligned}$$

as required. If $k \notin \sigma$ and $(\sigma \setminus \{l\}) \cup \{k\} \notin \nabla$, then $S_{kl}(\sigma) = (\sigma \setminus \{l\}) \cup \{k\}$. Hence $S_{kl}(\sigma) = ((\sigma_j \setminus \{l\}) \cup \{k\}) \cup \tau_1 = \sigma_j^* \cup \tau_1$, as required.

Case II) $S_{kl}(\sigma_j) = \sigma_j$ and there is no $s \neq j$ such that $\sigma_j = S_{kl}(\sigma_s) \cup \tau$. Then $\sigma_j^* = \sigma_j$. If $S_{kl}(\sigma) = \sigma$, then $S_{kl}(\sigma) = \sigma = \sigma_j^* \cup \tau_1$, as required. If $S_{kl}(\sigma) \neq \sigma$, then $l \in \sigma$, $k \notin \sigma$ and $(\sigma \setminus \{l\}) \cup \{k\} \notin \nabla$, hence $S_{kl}(\sigma) = (\sigma \setminus \{l\}) \cup \{k\} = ((\sigma_j \cup \tau_1) \setminus \{l\}) \cup \{k\}$. Note that $k \notin \sigma_j$ implies that $l \notin \sigma_j$, otherwise, from $S_{kl}(\sigma_j) = \sigma_j$ we have that $(\sigma_j \setminus \{l\}) \cup \{k\} \in \nabla$, then $((\sigma_j \cup \tau_1) \setminus \{l\}) \cup \{k\} = ((\sigma_j \setminus \{l\}) \cup \{k\}) \cup \tau_1 \in \nabla$, a contradiction. Hence $l \in \tau_1$ and

$$\begin{aligned} S_{kl}(\sigma) &= \sigma_j \cup ((\tau_1 \setminus \{l\}) \cup \{k\}) \\ &= \sigma_j^* \cup ((\tau_1 \setminus \{l\}) \cup \{k\}), \end{aligned}$$

as required.

Case III) $S_{kl}(\sigma_j) = \sigma_j$ and there is some $s \neq j$ such that $\sigma_j = S_{kl}(\sigma_s) \cup \tau$. Then $\sigma = \sigma_j \cup \tau_1 = S_{kl}(\sigma_s) \cup \tau \cup \tau_1$. Note that $S_{kl}(\sigma_s) \neq \sigma_s$, which implies that $k \in S_{kl}(\sigma_s) = (\sigma_s \setminus \{l\}) \cup \{k\}$. Then $k \in \sigma$ and $S_{kl}(\sigma) = \sigma$. Since $\sigma_s^* = S_{kl}(\sigma_s)$, it follows that $S_{kl}(\sigma) = \sigma = \sigma_s^* \cup (\tau \cup \tau_1)$, as required.

Secondly, we show that $x_{\sigma_1^*}, x_{\sigma_2^*}, \dots, x_{\sigma_r^*}$ is a minimal system. It is enough to show that $\sigma_i^* \not\subseteq \sigma_j^*$ for any $i \neq j$. Divide into three cases again.

Case 1) $\sigma_j^* = \sigma_j$. It is clear that $\sigma_i^* \not\subseteq \sigma_j^*$ if $\sigma_i^* = \sigma_i$ or $\sigma_t \cup \tau$, so we may assume that $\sigma_i^* = S_{kl}(\sigma_i)$. As $S_{kl}(\sigma_j) = \sigma_j$, if $\sigma_i^* \subseteq \sigma_j^*$ then $S_{kl}(\sigma_i) \subseteq S_{kl}(\sigma_j)$, which implies that $\sigma_j^* \neq \sigma_j$, a contradiction.

Case 2) $\sigma_j^* = S_{kl}(\sigma_j) \neq \sigma_j$. Then, as $S_{kl}(\sigma_j) \notin \nabla$, we see that $\sigma_i^* \not\subseteq \sigma_j^*$ if $\sigma_i^* = \sigma_i$ or $\sigma_t \cup \tau$. If $\sigma_i^* = S_{kl}(\sigma_i) \neq \sigma_i$ and $\sigma_i^* \subseteq \sigma_j^*$, then $l \in \sigma_i, \sigma_j$, $k \notin \sigma_i, \sigma_j$ and $(\sigma_i \setminus \{l\}) \cup \{k\} \subseteq (\sigma_j \setminus \{l\}) \cup \{k\}$. It follows that $\sigma_i \subseteq \sigma_j$, a contradiction.

Case 3) $\sigma_j^* = \sigma_t \cup \tau$, where $\sigma_t < \tau$. Then $\sigma_j = S_{kl}(\sigma_t) \cup \tau$, $l \in \sigma_t$, $k \notin \sigma_t$ and $S_{kl}(\sigma_t) = (\sigma_t \setminus \{l\}) \cup \{k\} \notin \nabla$. If $\sigma_i^* = S_{kl}(\sigma_i) \neq \sigma_i$, then $k \in \sigma_i^*$, hence $\sigma_i^* \not\subseteq \sigma_j^*$ as $k \notin \sigma_j^*$. If $\sigma_i^* = \sigma_s \cup \tau'$, where $\sigma_s < \tau'$, $l \in \sigma_s$ and $k \notin \sigma_s$, then $\sigma_i^* \subseteq \sigma_j^*$ implies that $\sigma_t \cup \tau = \sigma_s \cup \tau' \cup \tau''$ for some τ'' , it follows that $((\sigma_t \setminus \{l\}) \cup \{k\}) \cup \tau = ((\sigma_s \setminus \{l\}) \cup \{k\}) \cup \tau' \cup \tau''$, i.e., $\sigma_j = \sigma_i \cup \tau' \cup \tau''$, a contradiction. Finally, we assume that $\sigma_i^* = \sigma_i$ and $\sigma_i^* \subseteq \sigma_j^*$. Then $\sigma_t \cup \tau = \sigma_i \cup \tau_i$ for some $\tau_i \neq \emptyset$ and $S_{kl}(\sigma_i) = \sigma_i$. As $l \in \sigma_t$, $k \notin \sigma_t$ and $k < l$, we have that $l \in \sigma_i$ or $l \in \tau_i$, and $k \notin \sigma_i$ and $k \notin \tau_i$. If $l \in \sigma_i$, then, from $k \notin \sigma_i$ and $S_{kl}(\sigma_i) = \sigma_i$, we have that $(\sigma_i \setminus \{l\}) \cup \{k\} \in \nabla$. Thus $(\sigma_i \setminus \{l\}) \cup \{k\} = \sigma_u \cup \tau_u$ for some $\sigma_u \in G(I_\Delta)$. Then, from $\sigma_t \cup \tau = \sigma_i \cup \tau_i$, we have that

$$\begin{aligned} \sigma_j &= S_{kl}(\sigma_t) \cup \tau \\ &= ((\sigma_t \setminus \{l\}) \cup \{k\}) \cup \tau \\ &= ((\sigma_i \setminus \{l\}) \cup \{k\}) \cup \tau_i \\ &= \sigma_u \cup \tau_u \cup \tau_i, \end{aligned}$$

a contradiction. If $l \in \tau_i$, then, from $\sigma_t \cup \tau = \sigma_i \cup \tau_i$, we have that

$$\begin{aligned} \sigma_j &= S_{kl}(\sigma_t) \cup \tau \\ &= ((\sigma_t \setminus \{l\}) \cup \{k\}) \cup \tau \\ &= \sigma_i \cup ((\tau_i \setminus \{l\}) \cup \{k\}), \end{aligned}$$

a contradiction again. The proof is complete. $\square$

Now we can prove

Theorem 2.3. *If Δ is stable, then, for all i and j ,*

$$\beta_{i,j}(I_\Delta) = \beta_{i,j}(I_{\Delta^c}).$$

Proof. Suppose that $\Delta^c = \text{Shift}_{k_r l_r}(\text{Shift}_{k_{r-1} l_{r-1}}(\dots(\text{Shift}_{k_1 l_1}(\Delta))\dots))$, where $r \geq 1$ and $k_i < l_i$, $i = 1, 2, \dots, r$. By 2.1, $\text{Shift}_{k_i l_i}(\text{Shift}_{k_{i-1} l_{i-1}}(\dots(\text{Shift}_{k_1 l_1}(\Delta))\dots))$ is stable for $i = 1, 2, \dots, r$. Hence it is enough to show that $\beta_{i,j}(I_\Delta) = \beta_{i,j}(I_{\text{Shift}_{kl}(\Delta)})$ for any $1 \leq k < l \leq n$. Let $G(I_\Delta) = \{x_{\sigma_1}, x_{\sigma_2}, \dots, x_{\sigma_r}\}$, then, by 2.2, $G(I_{\text{Shift}_{kl}(\Delta)}) = \{x_{\sigma_1^*}, x_{\sigma_2^*}, \dots, x_{\sigma_r^*}\}$. Since I_Δ and $I_{\text{Shift}_{kl}(\Delta)}$ are squarefree stable and $m(\sigma_i^*) = m(\sigma_i)$,

$i = 1, 2, \dots, r$, it follows from the formula of Betti numbers of squarefree stable ideals that $\beta_{i,j}(I_\Delta) = \beta_{i,j}(I_{\text{Shift}_{kl}(\Delta)})$, as required. $\square$

Combining 1.5, 1.6 and 2.3, we get the following

Theorem 2.4. *If Δ is stable, then, for all i and j ,*

$$\beta_{i,j}(I_{\Delta^s}) = \beta_{i,j}(I_{\Delta^e}) = \beta_{i,j}(I_{\Delta^c}).$$

REFERENCES

- [1] A. Aramova, J. Herzog, and T. Hibi, Shifting operations and graded Betti Numbers, J. Algebr. Comb. 12(2000) 207-222.
- [2] A. Aramova, J. Herzog, and T. Hibi, Ideals with stable Betti numbers, Adv. Math. 152(2000) 72-77.
- [3] A. Aramova, J. Herzog, and T. Hibi, Squarefree lexsegment ideals, Math. Z. 228(1998) 353-378.
- [4] A. Aramova, J. Herzog, and T. Hibi, Gotzmann theorems for exterior algebras combinatorics, J. Algebra 191(1997) 174-211.
- [5] W. Bruns, J. Herzog, Cohen-Macaulay rings, Revised edition, Cambridge Studies in advanced mathematics 39, Cambridge University Press, 1998.
- [6] D. Eisenbud, Commutative Algebra with a View Towards Algebraic Geometry, Springer, 1995.
- [7] D. Eisenbud, A. Reeves, and B. Totaro, Initial ideals, Veronese subrings, and rates of algebras, Adv. Math. 109(1994) 168-187.
- [8] S. Eliahou, M. Kervaire, Minimal resolutions of some monomial ideals, J. Algebra 129(1990) 1-25.
- [9] J. Herzog, T. Hibi, Componentwise linear ideals, Nagoya Math. J. 153 (1999) 141-153.
- [10] R. P. Stanley, Combinatorics and Commutative Algebra, Birkhäuser, 1983.

Zhongming Tang
 Department of Mathematics
 Suzhou University
 Suzhou 215006
 P. R. China
 E-mail: zmtang@suda.edu.cn

Guifen Zhuang
 Department of Mathematics
 Suzhou University
 Suzhou 215006
 P. R. China